arXiv:0903.0238v2 [math.GT] 18 May 2009

SINGULAR SEIFERT SURFACES AND SMALE INVARIANTS FOR A FAMILY OF 3-SPHERE IMMERSIONS

TOBIAS EKHOLM AND MASAMICHI TAKASE

Abstract. A self-transverse immersion of the 2-sphere into 4-space with algebraic number of self intersection points equal to n induces an immersion of the circle bundle over the 2-sphere of Euler class $2n$ into 4-space. Precomposing the circle bundle immersions with their universal covering maps, we get for $n > 0$ immersions g_n of the 3-sphere into 4-space. In this note, we compute the Smale invariants of g_n. The computation is carried out by (partially) resolving the singularities of the natural singular map of the punctured complex projective plane which extends g_n.

As an application, we determine the classes represented by g_n in the cobordism group of immersions which is naturally identified with the stable 3-stem. It follows in particular that g_n represents a generator of the stable 3-stem if and only if n is divisible by 3.

1. Introduction

Consider a self-transverse immersion $f\colon S^2 \to \mathbb{R}^4$ of the 2-sphere S^2 into 4-space $\mathbb{R}^4$. Let $p \in \mathbb{R}^4$ be a self intersection point of f and assume that S^2 and $\mathbb{R}^4$ are oriented. Ordering the two local sheets of $f(S^2)$ intersecting at p we get an intersection number at p. Since the codimension is even this intersection number is independent of the ordering. We say that the sum of the intersection numbers over all the double points of f is its *algebraic number of double points*.

The normal bundle of an immersion $f\colon S^2 \to \mathbb{R}^4$ with algebraic number of double points equal to n is the 2-dimensional oriented 2-plane bundle ξ_{2n} of Euler number $2n$, see [30]. The unit sphere bundle of ξ_{2n} is the lens space $L(2n,1)$ and it follows by micro-extension that an immersion $f\colon S^2 \to \mathbb{R}^4$, with algebraically n double points, induces an immersion $f_{2n}\colon L(2n,1) \to \mathbb{R}^4$. Consider the universal ($2n$-fold) covering $\pi_{2n}\colon S^3 \to L(2n,1)$. Precomposing we get an immersion

$$g_n = f_{2n} \circ \pi_{2n}\colon S^3 \to \mathbb{R}^4. \tag{1.1}$$

Let $\mathrm{Imm}[S^m, \mathbb{R}^N]$ denote the group of regular homotopy classes of immersions $S^m \to \mathbb{R}^N$, where the group operation is induced by connected sum of immersions. The Smale-Hirsch h-principle implies that the group $\mathrm{Imm}[S^m, \mathbb{R}^N]$ is isomorphic to the m^{th} homotopy group $\pi_m(V_{N,m})$ of the Stiefel manifold $V_{N,m}$ of m frames in

---

2000 *Mathematics Subject Classification.* Primary 57R45; Secondary 57R42, 55Q45, 57M99, 57R90.

Key words and phrases. immersion; cobordism; 3-manifold; singular Seifert surface; stable homotopy of spheres; umbilic; singularity.

TE acknowledges support from the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine. MT is partially supported by the Grant-in-Aid for Young Scientists (B), MEXT, Japan.

$\mathbb{R}^N$. The isomorphism is given by the Smale invariant

$$\Omega\colon \operatorname{Imm}[S^m,\mathbb{R}^N] \to \pi_m(V_{N,m}),$$

see [25]. In particular, in the case studied in this paper:

$$\operatorname{Imm}[S^3,\mathbb{R}^4] = \pi_3(V_{4,3}) \approx \pi_3(SO_4) \approx \mathbb{Z}\oplus\mathbb{Z},$$

see Section 2.1 for a description of the isomorphism.

Let $\operatorname{Cob}(m,N)$ denote the cobordism group of immersions of oriented closed m-manifolds into $\mathbb{R}^N$, where the group operation is induced by disjoint union. By [21], the cobordism group $\operatorname{Cob}(3,4)$ is isomorphic to the stable 3-stem

$$\operatorname{Cob}(3,4) \approx \pi_3^S \approx \mathbb{Z}_{24}.$$

Theorem 1.1. *The Smale invariant of the immersion $g_n\colon S^3 \to \mathbb{R}^4$, $n>0$ satisfies*

$$\Omega(g_n) = \big(4n-1, -n(n+2)\big) \in \mathbb{Z}\oplus\mathbb{Z}.$$

It follows in particular that g_n represents the element

$$-(2n^2+1) \bmod 24 \in \mathbb{Z}_{24} \approx \pi_3^S$$

and hence generates the stable 3-stem if and only if n is a multiple of 3.

Theorem 1.1 is proved in Section 3. The proof uses singular Seifert surfaces, which were introduced in [5] and has since proved to be quite useful in the study of embeddings, immersions, and other maps. See for example [1, 5, 6, 9, 24, 29]. In fact, one of the main technical points of this paper is a concrete construction of a singular Seifert surfaces for the immersions g_n. The construction is based on perturbations derived from an unfolding of a certain complex map germ, regarded as a real map germ, and might be of interest from a more general viewpoint since similar constructions of stable- or other singular maps with desired properties can be used in many other settings.

In Section 4 we give a brief discussion of relations to other results. We first discuss the immersion $g_1\colon S^3 \to \mathbb{R}^4$, studied by Milnor [17, (11), §IV] and since then in several papers (e. g. [11, 4]). In this case, Theorem 1.1 recovers the results [4, Proposition 8.4.1] which shows that g_1 represents a generator in $\mathbb{Z}_8 \subset \mathbb{Z}_{24}$ as well as the fact that the triple point invariant $\lambda(g_1) \in \mathbb{Z}_3$ (defined in [4, §6.2]) of g_1 vanishes. Second, we show that the immersion g_2 coincides with Melikhov's example of an immersion with non-trivial stable Hopf invariant [16, Example 4].

Notational conventions. We work in the smooth category throughout; all manifolds and immersions are assumed to be differentiable of class C^∞. Furthermore, all spheres and Euclidean spaces are assumed to be oriented and we orient the boundary of an oriented manifold by the "outward normal first" convention, as in e.g. [8, 10].

2. SMALE INVARIANTS IN TERMS OF SINGULAR SEIFERT SURFACES

In this section we first establish notation for the Smale invariant of immersions $S^3 \to \mathbb{R}^4$ and review some of its properties. Then we present a formula for one of the components of the Smale invariant in terms of singularities of a Seifert surface.

2.1. **The regular homotopy class.** An immersion $f\colon S^3 \to \mathbb{R}^4$ comes equipped with a natural stable framing via the bundle isomorphism $\epsilon^1 \oplus TS^3 \cong f^*T\mathbb{R}^4$, where ϵ^1 denotes the trivial line bundle. The homotopy class of the stable framing is completely characterized by two integers: the degree $D(f)$ and the Hirzebruch defect $H(f)$. We refer to [10] for the detailed definitions and mention here only that $D(f)$ equals the normal degree of f, see [17].

The Smale-Hirsch h-principle implies that the map which associates to an immersion $S^3 \to \mathbb{R}^4$ its natural stable framing induces a (weak) homotopy equivalence between the space of immersions and the space of stable framings. In particular, on π_0 (i.e. on the level of path components) the Smale invariant, see Section 1,

$$\Omega(f) \in \pi_3(SO_4) \approx \mathbb{Z} \oplus \mathbb{Z},$$

which determines the regular homotopy class of an immersion $f\colon S^3 \to \mathbb{R}^4$, is determined by $D(f)$ and $H(f)$.

More precisely, we have the following expression for $\Omega\colon \mathrm{Imm}[S^3, \mathbb{R}^4] \to \mathbb{Z} \oplus \mathbb{Z}$,

$$\Omega([f]) = \left(D(f) - 1, \frac{H(f) - 2(D(f) - 1)}{4} \right) \in \mathbb{Z} \oplus \mathbb{Z}, \tag{2.1}$$

where $f\colon S^3 \to \mathbb{R}^4$ is a representative of $[f] \in \mathrm{Imm}[S^3, \mathbb{R}^4]$. To see this, use [10, Theorems 2.2(a) and 2.5(a)] which describe how $D(f)$ and $H(f)$ behave under the natural action of SO_4, expressed in terms of the two generators of $\pi_3(SO_4)$ given in [26, p. 117], in combination with [10, Theorems 2.2(b) and 2.5(b)] which show that $D(\iota) = 1$ and $H(\iota) = 0$ for the standard inclusion $\iota\colon S^3 \to \mathbb{R}^4$.

Formula (2.1) follows Hughes' convention for the Smale invariant [8, §2]. It is chosen so that for an immersion $f\colon S^3 \to \mathbb{R}^4$ which extends to an immersion $F\colon V \to \mathbb{R}^4$ of a compact oriented 4-manifold V with boundary $\partial V = S^3$, we get

$$\Omega(f) = \left(\chi(V) - 1, \frac{-3\sigma(V) - 2\chi(V) + 2}{4} \right) \in \mathbb{Z} \oplus \mathbb{Z},$$

see [8, Theorem 3.1], where $D(f) = \chi(V)$ is the Euler characteristic of V and $H(f) = -3\sigma(V)$ where $\sigma(V)$ is the signature of V, see [10, Theorems 2.2(b) and 2.5(b)].

Remark 2.1. Since $H(f) - 2D(f)$ is always congruent to 2 modulo 4 by [10, Theorem 2.6], the second component in the right-hand side of formula (2.1) is an integer.

2.2. **The Smale invariant in 5-space and cobordism class.** The long exact sequence of homotopy groups of the fibration

$$SO_2 \to SO_5 \to V_{5,3},$$

where $V_{5,3}$ is the Stiefel manifold of orthogonal 3-frames in $\mathbb{R}^5$, shows that $\pi_3(SO_5) = \pi_3(V_{5,3})$. We can thus consider the Smale invariant of an immersion $h\colon S^3 \to \mathbb{R}^5$ as an element

$$\Omega(h) \in \pi_3(SO_5) \approx \mathbb{Z}.$$

In particular if $f\colon S^3 \to \mathbb{R}^4$ is an immersion and if $j\colon \mathbb{R}^4 \to \mathbb{R}^5$ denotes the inclusion then the fact that the map $\pi_3(SO_4) \to \pi_3(SO_5)$ induced by the inclusion is given by $(a, b) \mapsto a + 2b$, see e.g. [8, p. 178], implies that the Smale invariant of $j \circ f$ satisfies

$$\Omega(j \circ f) = \frac{1}{2} H(f) \in \mathrm{Imm}[S^3, \mathbb{R}^5] \approx \mathbb{Z}.$$

As mentioned in Section 1, the oriented cobordism group $\mathrm{Cob}(3,4)$ of immersions of oriented 3-manifolds into $\mathbb{R}^4$ is isomorphic to the stable homotopy group $\pi_3^S \approx \mathbb{Z}_{24}$ of spheres. The isomorphism can be constructed via the Pontrjagin-Thom construction and in order to compute the cobordism class represented by an immersion $f\colon S^3 \to \mathbb{R}^4$ we use the following factorization of the map $\mathrm{Imm}[S^3, \mathbb{R}^4] \to \mathrm{Cob}(3,4)$:

$$\pi_3(SO_4) \longrightarrow \pi_3(SO_5) \longrightarrow \cdots \longrightarrow \pi_3(SO) \xrightarrow{J} \pi_3^S \approx \mathbb{Z}_{24},$$

where the last map is the J-homomorphism (see [8, p. 180]). In fact, we get an isomorphism

$$\mathrm{Cob}(3,4) \to \mathbb{Z}_{24}$$

by mapping an immersion $f\colon M \to \mathbb{R}^4$ representing a cobordism class of an immersion of a closed 3-manifold M to the element

$$\frac{H(f) + 3\mu(M, s_f)}{2} \pmod{24},$$

where $\mu(M, s_f) \in \mathbb{Z}_{16}$ is the μ-invariant of M with respect to the spin structure s_f induced by f, see [29]. Note that $\mu(S^3, s_f)$ is always equal to zero in $\mathbb{Z}_{16}$.

2.3. **Singular Seifert surfaces and Stingley's index.** For immersions $f\colon S^3 \to \mathbb{R}^4$ which do not extend to immersions of compact oriented 4-manifolds (as was assumed for the discussion in the last paragraphs of Section 2.1), the Hirzebruch defect can be read off from extensions by singular maps which are regular near the boundary. An extension by a smooth map with singularities of a given map (embedding, immersion, generic map, etc.) is often called a singular Seifert surface because of similarities with Seifert surfaces in knot theory. For immersions somewhat different definitions have been used after the notion was introduced in [5].

In [29], generic maps $F\colon V \to \mathbb{R}^4$ from a compact oriented 4-manifold V with $\partial V = S^3$, $F|_{\partial V} = f$, where $f\colon S^3 \to \mathbb{R}^4$ is an immersion, and with no singularities in a neighborhood of ∂V are considered. In this case, the Hirzebruch defect $H(f)$ is expressed as

$$H(f) = -3\sigma(V) - \sharp\Sigma^{2,0}(F)$$

where $\sharp\Sigma^{2,0}(F)$ is the (algebraic) number of umbilic points of F. This was essentially proved in [29, Proof of Lemma 3.2] (although the sign convention there is not the same as the one here; see Remark 2.6 below). The proof is based on the Thom polynomial for umbilic singularities of a generic map $G\colon M \to \mathbb{R}^4$ from a closed oriented 4-manifold M, see [13, 22] and also [2, §2,2 in Chapter 4]:

$$\sharp\Sigma^{2,0}(G) = -\langle p_1(M), [M]\rangle = -3\sigma(M), \tag{2.2}$$

where p_1 denotes the first Pontrjagin class and $[M] \in H_4(M;\mathbb{Z})$ the fundamental class.

For the purposes of this paper, we weaken the condition for a singular Seifert surfaces and use the following characterization.

Definition 2.2. Let $f\colon S^3 \to \mathbb{R}^4$ be an immersion. A *singular Seifert surface* for f is a smooth map $F\colon V \to \mathbb{R}^4$ from a compact oriented 4-manifold V with $\partial V = S^3$ which satisfies the following conditions. The restriction $F|_{\partial V}$ to the boundary equals f, there is a neighborhood of ∂V where the map F has no singularity, for any $p \in V$, the rank $\mathrm{rk}(dF_p)$ of the differential

$$dF_p\colon T_pV \to T_{F(p)}\mathbb{R}^4$$

satisfies $\mathrm{rk}(dF_p) \geq 2$, and points q with $\mathrm{rk}(dF_q) = 2$ are isolated.

Note that if $F\colon V \to \mathbb{R}^4$, $\partial V = S^3$ is a generic (i.e., locally stable) map which is non-singular near the boundary then its possible singularities are Morin singularities where the differential has rank 3 and isolated umbilic points where the differential has rank 2. Therefore a singular Seifert surface in the sense of [29, §2.1] fulfills the requirements of Definition 2.2.

Smooth maps between oriented 4-manifolds with differential of rank 2 at isolated points and > 2 elsewhere were studied by Stingley [27]. He associated an index [27, §2] to such isolated points as follows. (See Remark 2.6 below for details on sign conventions.) Let $g\colon M \to N$ be a smooth map between oriented 4-manifolds. Let $p \in M$ be such that $\mathrm{rk}(dg_p) = 2$ and assume that p is an isolated rank 2 point in the sense that there exists a neighborhood $U \subset M$ of p such that $\mathrm{rk}(dg_q) > 2$ for all $q \in U - \{p\}$. Then there are local coordinates $x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4$ around p and $y = (y_1, y_2, y_3, y_4) \in \mathbb{R}^4$ around $g(p)$ such that, in these coordinates the map $g(x) = \big(g_1(x), g_2(x), g_3(x), g_4(x)\big)$ is given by

$$\begin{cases} g_1(x) &= x_1,\\ g_2(x) &= x_2,\\ g_3(x) &= A(x),\\ g_4(x) &= B(x), \end{cases}$$

where $A(0) = B(0) = 0$ and where

$$\begin{pmatrix} \frac{\partial A}{\partial x_3} & \frac{\partial A}{\partial x_4} \\ \frac{\partial B}{\partial x_3} & \frac{\partial B}{\partial x_4} \end{pmatrix}$$

vanishes for $x = 0$ but not for $x \neq 0$.

Definition 2.3. The *index* $\mathrm{ind}_p(g)$ of an isolated rank 2 point p as above is

$$\mathrm{ind}_p(g) = \deg_\epsilon(\hat{g}),$$

where $\hat{g}(x) = \big(\hat{g}_1(x), \hat{g}_2(x), \hat{g}_3(x), \hat{g}_4(x)\big)$ is defined as

$$\begin{cases} \hat{g}_1(x) &= \frac{\partial A}{\partial x_3},\\ \hat{g}_2(x) &= \frac{\partial B}{\partial x_3},\\ \hat{g}_3(x) &= \frac{\partial A}{\partial x_4},\\ \hat{g}_4(x) &= \frac{\partial B}{\partial x_4}, \end{cases}$$

in a neighborhood of 0 and where $\deg_\epsilon(\hat{g})$ denotes the signed sum of preimages at any sufficiently small regular value ϵ of $\hat{g}$. (I.e. $\deg_\epsilon(\hat{g})$ is the degree of $\hat{g}$ around 0).

It is straightforward to check that $\mathrm{ind}_p(g)$ is independent of the choice of local coordinates. Furthermore, the index of an umbilic point of a map between 4-manifolds equals ± 1 and in this case we use the index to define the sign of umbilic points.

Remark 2.4. If the orientation of the source manifold in Definition 2.3 is changed then the index of each rank 2 point changes its sign. On the other hand, if the orientation of the target manifold is changed then the index of any rank 2 point remains unchanged.

For a smooth map $g\colon M \to N$ between oriented 4-manifolds where M is closed let

$$\Sigma^2(g) = \{p \in M;\ \mathrm{rk}(dg_p) = 2\}$$

and note that if the rank 2 points of g are isolated then $\Sigma^2(g)$ is finite. For such maps g with isolated rank 2 points we define *the algebraic number of rank* 2 *points* $\sharp\Sigma^2(g)$ of g as

$$\sharp\Sigma^2(g) = \sum_{p\in\Sigma^2(g)} \mathrm{ind}_p(g).$$

Using the above notation, there is the following Riemann-Hurwitz type formula for closed oriented 4-manifolds.

Theorem 2.5 (Stingley [27, Theorem I], see also Porter [18] and [7, §8])**.** *Let* $g\colon M \to N$ *be a smooth map between oriented* 4*-manifolds with isolated rank* 2 *points and assume that* M *is closed. Then*

$$\big\langle (g^*p_1(N) - p_1(M)), [M]\big\rangle = \sharp\Sigma^2(g),$$

where p_1 *denotes the first Pontrjagin class and* $[M]$ *is the fundamental class of* M*.*

In particular, Theorem 2.5 implies that if $g\colon M \to \mathbb{R}^4$ is a map from a closed oriented 4-manifold M with isolated rank 2 points, then

$$\sharp\Sigma^2(g) = -\big\langle p_1(M), [M]\big\rangle = -3\sigma(M), \tag{2.3}$$

where $\sigma(M)$ is the signature of M.

Remark 2.6. The Thom polynomial (2.2) and Stingley's formula (2.3) imply that for an appropriate choice of signs, the count of indices of rank 2 points of maps of a closed 4-manifold M into $\mathbb{R}^4$ gives the first Pontrjagin number of M. However, the sign conventions for the index which appear in the literature are often conflicting (see [13, 18, 22, 23, 27, 29] and also [2, §2.2 in Chapter 4], for example). Even within Stingley's paper [27], signs appear to be in conflict: compare the signs in [27, Lemma 2.1] to those in the first line of the proof of [27, Lemma 4.1].

Definition 2.3 above adopts the sign convention of [27, Lemma 4.1], which is opposite to [27, Lemma 2.1]. The computations in the proof of Theorem 1.1 below, together with Remark 2.1, show that the sign convention of Definition 2.3 is a reasonable choice.

2.4. **A formula for the Hirzebruch defect.**

Let $f\colon S^3 \to \mathbb{R}^4$ be an immersion and let $F\colon V \to \mathbb{R}^4$ be a singular Seifert surface of f, see Definition 2.2.

Proposition 2.7. *The Hirzebruch defect* $H(f)$ *(of the stable framing induced by* f*) satisfies*

$$H(f) = -3\sigma(V) - \sharp\Sigma^2(F). \tag{2.4}$$

In particular, the right hand side of (2.4) *does not depend on the choice of singular Seifert surface* $F\colon V \to \mathbb{R}^4$ *extending* f*.*

Proof. Let $F'\colon V' \to \mathbb{R}^4$ be a singular Seifert surface for an immersion $f'\colon S^3 \to \mathbb{R}^4$ which is regularly homotopic to f. Using F', F, and the trace of a regular homotopy $h\colon S^3 \times [0,1] \to \mathbb{R}^4 \times [0,1]$ between f and f', we construct a map

$$\hat{G}\colon V \cup_\partial S^3 \times [0,1] \cup_\partial (-V') \to \mathbb{R}^4 \times [0,1]$$

from the closed oriented 4-manifold $W = V \cup_\partial S^3 \times [0,1] \cup_\partial (-V')$, where $-V'$ denotes the manifold V' with reversed orientation, which is smooth after smoothing corners. Composing $\hat{G}$ with a generic projection to $\mathbb{R}^4$, we obtain a smooth map $G\colon W \to \mathbb{R}^4$ with stable singularities, see [15]. In particular, its rank 2 points are isolated.

The number of rank 2 points $\sharp\Sigma^2(G)$ satisfies

$$\sharp\Sigma^2(G) = \sharp\Sigma^2(F) - \sharp\Sigma^2(F'),$$

since the trace of a regular homotopy h is an immersion which after composition with a projection does not have any rank 2 points.

By (2.3), we have $\sharp\Sigma^2(G) = -3\sigma(W)$ and hence

$$-3\sigma(V) - \sharp\Sigma^2(F) = -3\sigma(V') - \sharp\Sigma^2(F')$$

by Novikov additivity. It follows that

$$-3\sigma(V) - \sharp\Sigma^2(F)$$

depends only on the regular homotopy class of f. We can thus define a homomorphism $a\colon \operatorname{Imm}[S^3, \mathbb{R}^4] \to \mathbb{Z}$ as follows. For $[f] \in \operatorname{Imm}[S^3, \mathbb{R}^4]$, pick a representative immersion f of $[f]$ and a Seifert surface $F\colon V \to \mathbb{R}^4$ of f and define

$$a([f]) = -3\sigma(V) - \sharp\Sigma^2(F).$$

Consider the subgroup $E \subset \operatorname{Imm}[S^3, \mathbb{R}^4]$ corresponding to

$$E = \big\{[f] \in \operatorname{Imm}[S^3, \mathbb{R}^4];\ \Omega(f) = (m,n) \in \mathbb{Z} \oplus \mathbb{Z},\ m + 2n \in 24\mathbb{Z}\big\},$$

where f denotes a representative for the class $[f]$ and where Ω is the Smale invariant. It follows from the discussion in Section 2.2 that

$$E = \ker\big(\operatorname{Imm}[S^3, \mathbb{R}^4] \to \operatorname{Cob}(3,4)\big),$$

see [8, Theorem 3.1]. In other words, each class of E corresponds to the class of an immersion which admits an immersed Seifert surface. By [10, Theorems 2.5(b)], the homomorphism a coincides with the Hirzebruch defect on E and hence the two homomorphisms agree on all of $\operatorname{Imm}[S^3, \mathbb{R}^4]$. □

3. Explicit constructions of singular Seifert surfaces

In this section we first construct singular Seifert surfaces for the immersions $S^3 \to \mathbb{R}^4$ described in Section 1 and then use these to prove Theorem 1.1.

We will use the following notation throughout this section. The 2-plane bundle over S^2 of Euler number k will be denoted ξ_k, $E(\xi_k)$ will denote the total space of the corresponding 2-disk bundle, and $L(k,1)$ the total space of the corresponding circle bundle ($L(k,1)$ is a $(k,1)$-lens space). Furthermore, recall from Section 1 that the normal bundle of a self-transverse immersion $f\colon S^2 \to \mathbb{R}^4$ with algebraic number of double points equal to n is ξ_{2n}. Thus, micro-extension gives an immersion $F_{2n}\colon E(\xi_{2n}) \to \mathbb{R}^4$ which agrees with f when restricted to the 0-section. Precomposing with the universal covering $\pi_{2n}\colon S^3 \to L(2n,1) = \partial E(\xi_{2n})$, we get the immersions

$$g_n = F_{2n} \circ \pi_{2n}\colon S^3 \to \mathbb{R}^4, \quad n > 0.$$

We next construct natural fillings of the maps g_n. To this end, note that the space $L(1,1) = \partial E(\xi_1)$ is S^3, thought of as the total space of the Hopf fibration, and that $E(\xi_1)$ is $\mathbb{C}P^2_0$, the complement of an open disk in the complex projective plane. Consider the k-fold branched cover

$$\Pi_k\colon E(\xi_1) \to E(\xi_k)$$

which extends the universal cover

$$\pi_k\colon S^3 = \partial E(\xi_1) \to \partial E(\xi_k) = L(k,1),$$

and which is the k-fold branched cover with a single branch point at the origin in each fiber disk. The map $G_n\colon \mathbb{C}P^2_0 \to \mathbb{R}^4$,

$$G_n = F_{2n} \circ \Pi_{2n}$$

then extends the immersion $g_n\colon S^3 \to \mathbb{R}^4$ but is *not* a singular Seifert surface for g_n for the following reason. The rank of dG_n equals 4 outside the 0-section in $E(\xi_1)$ (in $\mathbb{C}P^2_0 - \mathbb{C}P^1$) and equals 2 at any point on the 0-section (along $\mathbb{C}P^1 \subset \mathbb{C}P^2_0$) and thus its rank 2 points are not isolated.

The rank 2 points of G_n forming a submanifold is reminiscent of the critical locus of a Morse-Bott function. In analogy with the Morse-Bott case, we will construct a perturbation of G_n below which creates a map with exactly two isolated rank 2 points. Furthermore, we will compute the indices of these rank 2 points. In fact, our perturbation of G_n has the form $F_{2n} \circ \Pi^{\epsilon}_{2n}$ where Π^{ϵ}_{2n} is a perturbation of the branched cover Π_{2n}. In order to construct the perturbation, we use the following local coordinate description of the bundles ξ_k.

Consider the decomposition of S^2 into two disks, the northern- and southern hemispheres D_N and D_S, respectively. After identification of these disks with the unit disk $D = \{w \in \mathbb{C};\ |w| \leq 1\}$ in the complex plane $\mathbb{C}$, we have

$$S^2 = D_S \cup_\psi D_N,$$

where $\psi\colon\ \partial D_S \to \partial D_N$ is the map given by $\psi(e^{i\theta}) = e^{-i\theta}$. The total spaces $E(\xi_k)$ can then be described as follows:

$$E(\xi_k) = D_S \times D \cup_{\psi_k} D_N \times D,$$

where $\psi_k\colon \partial D_S \times D \to \partial D_N \times D$ is given by

$$\psi_k(e^{i\theta}, z) = (e^{-i\theta}, e^{-ki\theta} z).$$

To see this, it is sufficient to note that the clutching function described gives a bundle with first Chern class (Euler number) equal to k.

In these local coordinates the map $\Pi_k\colon E(\xi_1) \to E(\xi_k)$ is the following:

$$\Pi_k(w,z) = \begin{cases} (w, z^k) \in D_S \times D & \text{if } (w,z) \in D_S \times D, \\ (w, z^k) \in D_N \times D & \text{if } (w,z) \in D_N \times D. \end{cases}$$

To see that this formula defines a map as claimed, note that on the overlap $(e^{i\theta}, z) \in \partial D_S \times D \subset E(\xi_1)$ is identified with $(e^{-i\theta}, e^{-i\theta} z) \in \partial D_N \times D \subset E(\xi_1)$. The former maps to $(e^{i\theta}, z^k) \in \partial D_S \times D \subset E(\xi_k)$ which is identified with $(e^{-i\theta}, e^{-ki\theta} z^k) \in \partial D_N \times D \subset E(\xi_k)$ which in turn is the image of the latter.

In order to perturb $\Pi_k\colon E(\xi_1) \to E(\xi_k)$ we consider the restriction of this map to fibers: $h(z) = z^k$. Let c be any complex number and let $h_c(z) = z^k + \bar{z}c$.

Lemma 3.1. *The singular set of h_c is the circle*

$$\Sigma = \bigl\{z;\ |z| = (k^{-1}|c|)^{\frac{1}{k-1}}\bigr\}.$$

If $c \neq 0$*, then the rank of* dh_c *equals* 1 *along* Σ *and the kernel field* $\ker(dh_c)$ *is transverse to* Σ *except at* $k+1$ *cusp points where the kernel field has simple tangencies with* Σ*. In particular,* h_c *is a stable map for* $c \neq 0$*.*

Proof. If ξ is a tangent vector to $\mathbb{C}$ thought of as a complex number we have

$$dh_c(\xi) = kz^{k-1}\xi + c\bar{\xi}.$$

Consequently, dh_c has kernel at points z where the equation $kz^{k-1}\xi + c\bar{\xi} = 0$ has non-trivial solutions. This is the case if

$$kz^{k-1} = c\bar{\xi}/\xi, \tag{3.1}$$

or in other words when $z \in \Sigma$. The properties of the kernel field are straightforward consequences of (3.1). □

We next deform the map Π_k using the perturbation described above on fibers. Note however that c in the above discussion must then be replaced by the image of a section of $E(\xi_1)$ and hence we cannot avoid rank 2 points.

Fix $\epsilon \in (0, \frac{1}{4})$. Define $\Pi_k^\epsilon \colon E(\xi_1) \to E(\xi_k)$ as follows, using the local coordinates introduced above,

$$\Pi_k^\epsilon(w,z) = \begin{cases} \big(w, (1-\epsilon)z^k + \epsilon w\bar{z}\big) \in D_S \times D & \text{if } (w,z) \in D_S \times D, \\ \big(w, (1-\epsilon)z^k + \epsilon w^k\bar{z}\big) \in D_N \times D & \text{if } (w,z) \in D_N \times D. \end{cases} \tag{3.2}$$

To see that (3.2) indeed defines a map, note that for $(w,z) = (e^{i\theta}, z) \in \partial D_S \times D$,

$$\psi_k\left(e^{i\theta}, (1-\epsilon)z^k + \epsilon e^{i\theta}\bar{z}\right) = \left(e^{-i\theta}, (1-\epsilon)e^{-ik\theta}z^k + \epsilon e^{-i(k-1)\theta}\bar{z}\right) \in \partial D_N \times D,$$

which is the image of the corresponding point $\psi_1(w,z) = (e^{-i\theta}, e^{-i\theta}z) \in \partial D_N \times D$ according to the second row in the right hand side of (3.2). Furthermore if $|z| \leq 1$ and $|w| \leq 1$ then $|(1-\epsilon)z^k + \epsilon w\bar{z}| \leq 1$ so that second components in the vectors in the right hand side of (3.2) lies in D.

Lemma 3.2. *The map* $\Pi_k^\epsilon \colon \mathbb{C}P_0^2 \to E(\xi_k)$ *is a map with two isolated rank 2 points:* p_S *corresponding to* $(0,0) \in D_S \times D$ *and* p_N *corresponding to* $(0,0) \in D_N \times D$*. Furthermore,* $\mathrm{ind}_{p_S}(\Pi_k^\epsilon) = k-1$ *and* $\mathrm{ind}_{p_N}(\Pi_k^\epsilon) = k(k-1)$*.*

Proof. If a tangent vector to $D_S \times D$ or $D_N \times D$ is viewed as a pair of complex numbers (η, ξ) and if $\epsilon' = \frac{\epsilon}{1-\epsilon}$ then, with h_c as in Lemma 3.1,

$$\big[d\Pi_k^\epsilon(w,z)\big](\eta,\xi) =$$
$$\begin{cases} \big(\eta, (1-\epsilon)\big[dh_{\epsilon' w}(z)\big](\xi) + \epsilon\bar{z}\eta\big) \in T(D_S \times D) & \text{if } (w,z) \in D_S \times D, \\ \big(\eta, (1-\epsilon)\big[dh_{\epsilon' w^k}(z)\big](\xi) + \epsilon k w^{k-1}\bar{z}\eta\big) \in T(D_N \times D) & \text{if } (w,z) \in D_N \times D. \end{cases}$$

Lemma 3.1 then implies that p_S and p_N are the only rank 2 points of Π_k^ϵ.

To compute the indices we consider the map $H \colon \mathbb{C}^2 \to \mathbb{C}^2$

$$H(z,w) = \left(w, z^\ell + \bar{z}w^m\right),$$

where $\ell \geq 2$ and $m \geq 1$. Clearly the only rank 2 point of H is the origin. To compute the index $\mathrm{ind}_0(H)$, write $K(w,z) = z^\ell + \bar{z}w^m$, then according to Definition 2.3, the index equals the degree near 0 of the map $\kappa \colon \mathbb{C}^2 \to \mathbb{C}^2$ where

$$\kappa(w,z) = \left(\frac{\partial K}{\partial z} + \frac{\partial K}{\partial \bar{z}},\ i\left(\frac{\partial K}{\partial z} - \frac{\partial K}{\partial \bar{z}}\right)\right) = \left(\ell z^{\ell-1} + w^m, i(\ell z^{\ell-1} - w^m)\right).$$

Note that there exists a complex linear isomorphism L of $\mathbb{C}^2$ such that

$$L \circ \kappa(w,z) = \left(w^m, z^{\ell-1}\right).$$

Since κ is holomorphic, the local degree at any inverse image of a regular value equals $+1$. Hence if ϵ is any sufficiently small regular value of κ we have

$$\operatorname{ind}_0(H) = \deg_\epsilon(\kappa) = m(\ell-1).$$

Taking $(m,\ell) = (k,1)$ and $(m,\ell) = (k,k)$, the statements on the indices of p_S respectively p_N follow. □

Proof of Theorem 1.1. Fix a small $\epsilon > 0$ and write $\pi_k^\epsilon = \Pi_k^\epsilon|_{S^3}$, $S^3 = \partial E(\xi_1)$ and note that

$$\pi_k^{s\epsilon} \colon S^3 \to E(\xi_k), \quad 0 \le s \le 1$$

gives a regular homotopy connecting the immersions $\pi_k \colon S^3 \to L(k,1) \subset E(\xi_k)$ to $\pi_k^\epsilon \colon S^3 \to E(\xi_k)$. Consequently the map

$$g_n^\epsilon = F_{2n} \circ \pi_{2n}^\epsilon \colon S^3 \to \mathbb{R}^4$$

is regularly homotopic to $g_n \colon S^3 \to \mathbb{R}^4$. Furthermore, since $F_{2n} \colon E(\xi_{2n}) \to \mathbb{R}^4$ is an immersion, Lemma 3.2 implies that

$$G_n^\epsilon = F_{2n} \circ \Pi_{2n}^\epsilon \colon \mathbb{C}P_0^2 \to \mathbb{R}^4$$

is a singular Seifert surface of g_n^ϵ with

$$\sharp\Sigma^2(G_n^\epsilon) = 2n - 1 + 2n(2n-1) = 4n^2 - 1.$$

Proposition 2.7 then gives the Hirzebruch defect

$$H(g_n) = H(g_n^\epsilon) = -3\sigma(\mathbb{C}P_0^2) - 4n^2 + 1 = -4n^2 - 2.$$

By [10, Theorem 2.2(b)], the normal degree of the immersion $f_{2n} \colon E(\xi_{2n}) \to \mathbb{R}^4$ equals $\chi(E(\xi_{2n})) = \chi(S^2) = 2$. Since $\pi_{2n} \colon S^3 \to L(2n,1)$ has degree $2n$, the composition $g_n = f_{2n} \circ \pi_{2n}$ has normal degree

$$D(g_n) = 4n$$

(see also [17, §IV]). Therefore, by the formula (2.1), we have

$$\Omega(g_n) = \left(4n-1, \frac{-4n^2 - 2 - 2\cdot(4n-1)}{4}\right) = (4n-1, -n(n+2)).$$

With this established, remaining statements are immediate consequences of the formulas in Section 2.2. □

4. Relations to other results

The immersion g_1 was apparently first studied by Milnor [17, (11), §IV] and has since appeared in several papers (e. g. [11, 4]). It was shown in [4, Proposition 8.4.1] that g_1 has odd Brown invariant (in $\mathbb{Z}_8$) and hence represents a generator in $\mathbb{Z}_8$ (see also [28, Lemma 1.7(a)] and [29, Remark 3.6]). Theorem 1.1 implies this result and furthermore that the triple point invariant $\lambda(g_1) \in \mathbb{Z}_3$ of g_1 vanishes. For a self-transverse immersion regularly homotopic to $j \circ g_1 \colon S^3 \to \mathbb{R}^4 \hookrightarrow \mathbb{R}^5$, the triple point invariant is defined to be the linking number modulo 3 of its image and a push-off of its double point set. For details, see [4, §6.2], and also [4, §9.1, p. 189] (where the letter "τ" should be replaced by "λ").

Melikhov studied in [16, Example 4] a construction similar to the one above. More precisely, he looked at the composition $e \circ q\colon S^3 \to \mathbb{R}^4$ of the universal 8-fold covering $q\colon S^3 \to Q^3$ and an embedding $e\colon Q^3 \to \mathbb{R}^4$ of the quaternion space $Q^3 = S^3/\{\pm 1, \pm i, \pm j, \pm k\}$ and showed that it represents an element with non-trivial stable Hopf invariant in the stable 3-stem $\pi_3^S \approx \mathbb{Z}_{24}$. Here, we show that his immersion essentially coincides with g_2.

Proposition 4.1. *The immersion $e \circ q\colon S^3 \to \mathbb{R}^4$ coincides with g_2 (up to orientation).*

Proof. Let $\hat{e}\colon \mathbb{R}P^2 \hookrightarrow \mathbb{R}^4$ be an embedding. Then, by [14, §2], the tubular neighborhood $N(\mathbb{R}P^2)$ of $\hat{e}(\mathbb{R}P^2) \subset \mathbb{R}^4$ is diffeomorphic to the total space of the non-orientable D^2-bundle over $\mathbb{R}P^2$ with Euler number ± 2 (see also [3, 19, 20, 31]). Furthermore, the boundary $\partial N(\mathbb{R}P^2)$ is known to be diffeomorphic to the quaternion space Q^3 (see [19]). We thus obtain an embedding $e\colon Q^3 \hookrightarrow \mathbb{R}^4$.

For the 2-fold covering $\rho\colon S^2 \to \mathbb{R}P^2$ and a suitable choice of the embedding $\hat{e}\colon \mathbb{R}P^2 \hookrightarrow \mathbb{R}^4$, the composition $\hat{e} \circ \rho\colon S^2 \to \mathbb{R}^4$ becomes an immersion with normal Euler class 4. Furthermore, if we denote by Γ the quaternion group $\{\pm 1, \pm i, \pm j, \pm k\}$ of order 8 and put $\mathbb{Z}_4 = \langle g;\ g^4 = 1\rangle$, then from the sequence

$$\{1\} \longrightarrow \mathbb{Z}_2 \longrightarrow \Gamma \longrightarrow \mathbb{Z}_4 \longrightarrow \{1\},$$

we have the sequence of the coverings

$$S^3 \longrightarrow L(4,1) = S^3/\mathbb{Z}_4 \longrightarrow Q^3 = S^3/\Gamma,$$

up to orientation [32]. Hence, we obtain the diagram

$$\begin{tikzcd}
S^3 \arrow[r, "\text{4-fold}"] \arrow[dr, "\text{8-fold}"'] & L(4,1) \arrow[r, "\text{Euler class 4}"'] \arrow[d, "\text{2-fold}"'] & S^2 \arrow[d, "\rho"'] \arrow[dr, "f_4|_{\partial E(\xi_4)}"] & \\
& Q^3 \arrow[r, "\text{Euler number 2}"'] & \mathbb{R}P^2 \arrow[r, "\hat{e}"'] & \mathbb{R}^4,
\end{tikzcd}$$

which is commutative up to orientation, see [12]. The proposition follows. ☐

Acknowledgment

MT would like to thank Professor Yuichi Yamada for illustrating various descriptions of the quaternion space and for valuable discussions, and Professor Takashi Nishimura for instructive discussions, especially on singularities of smooth mappings.

References

[1] Andersson, O.: A geometric classification of the path components of the space of locally stable maps $S^3 \to \mathbb{R}^4$, *Topology* **46** (2007), 577–597.

[2] Arnol'd, V. I., Vasil'ev, V. A., Goryunov, V. V. and Lyashko, O. V.: Singularities, Local and global theory, Translation by A. Iacob. Translation edited by V. I. Arnol'd. *Encyclopaedia of Mathematical Sciences* **6**, Dynamical Systems VI, Springer-Verlag, Berlin, 1993.

[3] Banchoff, T. F. and Johnson, O.: The normal Euler class and singularities of projections for polyhedral surfaces in 4-space, *Topology* **37** (1998), 419–439.

[4] Ekholm, T.: Differential 3-knots in 5-space with and without self-intersections, *Topology* **40** (2001), 157–196.

[5] Ekholm, T. and Szűcs, A.: Geometric formulas for Smale invariants of codimension two immersions, *Topology* **42** (2003), 171–196.

[6] Ekholm, T. and Szűcs, A.: The group of immersions of homotopy $(4k-1)$-spheres, *Bull. London Math. Soc.* **38** (2006), 163–176.

[7] Harvey, R. and Lawson, H. B.: Geometric residue theorems, *Amer. J. Math.* **117** (1995), 829–873.

[8] Hughes, J.: Bordism and regular homotopy of low-dimensional immersions, *Pacific J. of Math.* **156** (1992), 155–184.

[9] Juhász, A.: A geometric classification of immersions of 3-manifolds into 5-space, *Manuscripta Math.* **117** (2005), 65–83.

[10] Kirby, R. and Melvin, P.: Canonical framings for 3-manifolds, Proceedings of 6th Gökova Geometry-Topology Conference. *Turkish J. Math.* **23** (1999), 89–115.

[11] Lashof, R. and Smale, S.: On the immersion of manifolds in euclidean space, *Ann. of Math. (2)* **68** (1958), 562–583.

[12] Lawson, T.: Splitting S^4 on $\mathbb{R}P^2$ via the branched cover of $\mathbb{C}P^2$ over S^4, *Proc. Amer. Math. Soc.* **86** (1982), 328–330.

[13] MacPherson, R.: Singularities of vector bundle maps, Proceedings of Liverpool Singularities–Symposium, I (1969/70), 316–318, *Lecture Notes in Math.* **192**, Springer, Berlin, 1971.

[14] Massey, W. S.: Proof of a conjecture of Whitney, *Pacific J. of Math.* **31** (1969), 143–156.

[15] Mather, J.: Generic projections, *Ann. of Math. (2)* **98** (1973) 226–245.

[16] Melikhov, S. A.: Sphere eversions and the realization of mappings, *Tr. Mat. Inst. Steklova* **247** (2004), *Geom. Topol. i Teor. Mnozh.*, 159–181; translation in *Proc. Steklov Inst. Math.* 2004, no. 4 (247), 143–163.

[17] Milnor, J.: On the immersion of n-manifolds in $(n+1)$-space, *Comment. Math. Helv.* **30** (1956), 275–284.

[18] Porter, R.: Characteristic classes and singularities of mappings, Differential geometry (Proc. Sympos. Pure Math., Vol. XXVII, Part 1, Stanford Univ., Stanford, Calif., 1973), pp. 397–402. Amer. Math. Soc., Providence, R. I., 1975.

[19] Price, T. M.: Homeomorphisms of quaternion space and projective planes in four space, *J. Austral. Math. Soc. Ser. A* **23**, (1977) 112–128.

[20] Price, T. M. and Roseman, D. M.: Embeddings of the projective plane in four space, preprint (1975).

[21] Rohlin, V. A.: New results in the theory of four-dimensional manifolds (Russian), *Doklady Akad. Nauk SSSR (N.S.)* **84** (1952), 221–224; French translation by S. Ochanine: 'Quatre articles de V. A. Rohlin', À la recherche de la topologie perdue, ed. L. Guillou and A. Marin, Progr. Math. 62 (Birkhäuser, Boston, MA, 1986) 17–23.

[22] Ronga, F.: Le calcul de la classe de cohomologie entière duale à $\overline{\Sigma}^k$, Proceedings of Liverpool Singularities–Symposium, I (1969/70), 313–315, *Lecture Notes in Math.* **192**, Springer, Berlin, 1971.

[23] Saeki, O. and Sakuma, K.: Stable maps between 4-manifolds and elimination of their singularities, *J. London Math. Soc.* **59** (1999), 1117–1133.

[24] Saeki, O., Szűcs, A. and Takase, M.: Regular homotopy classes of immersions of 3-manifolds into 5-space, *Manuscripta Math.* **108** (2002), 13–32.

[25] Smale, S.: The classification of immersions of spheres in Euclidean spaces, *Ann. of Math. (2)* **69** (1959) 327–344.

[26] Steenrod, N.: The Topology of Fibre Bundles, Princeton Mathematical Series **14**, Princeton University Press, Princeton, N. J., 1951.

[27] Stingley, R.: Singularities of maps between 4-manifolds, dissertation, State University of New York at Stony Brook, 1995.

[28] Szűcs, A.: Two theorems of Rokhlin, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **267** (2000), *Geom. i Topol.* **5** 274–281, 332; translation in *J. Math. Sci.* (N. Y.) **113** (2003) 888–892.

[29] Takase, M.: An Ekholm–Szűcs-type formula for codimension one immersions of 3-manifolds up to bordism, *Bull. London Math. Soc.* **32** (2007) 163–178.

[30] Whitney, H.: The self-intersections of a smooth n-manifold in $2n$-space, *Ann. of Math. (2)* **45** (1944) 220–246.

[31] Yamada, Y.: Decomposition of S^4 as a twisted double of a certain manifold, *Tokyo J. Math.* **20** (1997) 23–33.

[32] Yamada, Y.: On a recognition of quaternionic space, private notes.

Department of mathematics, Uppsala University, Box 480, 751 06 Uppsala, Sweden
E-mail address: `tobias@math.uu.se`

Department of Mathematical Sciences, Faculty of Science, Shinshu University, Matsumoto, 390-8621 Japan.
E-mail address: `takase@math.shinshu-u.ac.jp`